



TOXICITY OF MERCURY ($HgCl_2$) ON CATFISH, *Clarias gariepinus*

Quazi Saleem^{1*}, Seema Hashmi²

¹Dept of Zoology, Milliya college Beed

² Dept. of Microbiology, Milliya college Beed

*Corresponding author email seemaseemaib@gmail.com

Abstract: In order to assess the acute toxicity of mercury on *Clarias gariepinus*, fish having mean weight and length of $32\text{ g} \pm 1.82$ and $18.2\text{cm} \pm 0.61$ were divided into six groups of six fish each. The different groups were exposed to the different concentrations of 0.5 mg/L, 1.0 mg/L, 1.5.8 mg/L, 2 mg/L 2.5 mg/L & 3.0mg/L for a period of 96 hours. One contrpl was also kept without Hgcl2. The results revealed that all the fish of groups exposed to 2.5, 3.0 & 3.5 mg/L died. The determination of 96 hours LC50 was carried out by computing the mortality results in Probit program of SPSS (version 17.0).

Keywords: *Clarias gariepinus*, mercury, acute toxicity, lethal concentration.

INTRODUCTION

In modern times, one of the main threats to the health of ecosystems is the exposure to a myriad of toxic substances and compounds such as mercury, cadmium, lead, copper, arsenic, air pollutants, pesticides, plastics, cigarette smoke, diesel fumes and nano-particles found in products like perfumes and sunscreens. Because of their high toxicity conferred by their persistent nature in the environment, heavy metals come to the forefront of dangerous substances causing serious health hazard in ecosystems and organisms[1-3]. Their introduction into aquatic environment is caused by direct or indirect agricultural and industrial discharges. Heavy metals contamination could be detrimental to ecological balance of the recipient environment and to a diversity of aquatic organisms [4 ,5]. Since fish are animals particularly affected by these pollutants they are widely used to evaluate the health of aquatic ecosystems[6].

Some particular heavy metals, such as mercury (Hg), are especially of a deep concern due to their high toxicity. Mercury occurs naturally as a mineral and is widely distributed throughout the environment as a result of natural and human activities. Inorganic mercury is the most common form of the metal released by industries in the environment [7]. Once in the aquatic ecosystem, part of the inorganic mercury can be

microbiologically converted into methyl-mercury.

The resulting organomercury compounds are rapidly absorbed by the gastrointestinal tract where 90% of ingested mercury is directly absorbed by fish from water through gills, skin and digestive tract[8]. Once contaminated by Hg, fish suffer pathological alterations, with consequent inhibition of metabolic processes, haematological changes, and decline in fertility and survival [9].

The aim of this study was to determine the acute toxicity of mercury chloride to *Clarias gariepinus*. *C gariepinus* was chosen as an experimental animal on the basis of important criteria. The fish is a representative species and widely used in aquaculture throughout India . Moreover it is largely tolerant of high concentration of heavy metals. In addition, *Clarias gariepinus* is a hardy fish and can survive difficult conditions.

MATERIAL AND METHODS

one hundred and fifty juveniles of *Clarias gariepinus* of the same brood were purchased from a reputable fish farm and transported in an oxygen bag to the aquaculture laboratory. Their mean weight and length were $32\text{ g} \pm 1.82$ and $18.2\text{cm} \pm 0.61$ respectively. The fish were kept in plastic tanks (30x 30x60cm) which were half filled with dechlorinated water. They were acclimatized to laboratory conditions over six weeks. During acclimatization, the juveniles were fed with

commercial fish feed known as Coppens at 4% of their body weight. The Juveniles were fed thrice daily (morning, afternoon and evening) and the water was changed every twenty four hours to prevent the accumulation of waste metabolite and food particles. They were kept at 12 hours of photoperiod. During the whole period of acclimatization, the mortality recorded was below 2%. At the end of acclimatization the new mean weight and length were $51.27g \pm 2.01$ and $20.2\text{ cm} \pm 0.72$ respectively. The juveniles were randomly divided into six groups of Eighteen fish and each group subdivided into a set of three subgroups consisting of six fish each. The eighteen subgroups were kept in eighteen different tanks in 15 liters of water. The six groups were respectively exposed to 0.5 mg/L, 1.0 mg/L, 1.5 mg/L, 2.0mg/L, 2.5mg/L and 3 mg/L of mercury chloride (HgCl₂) one control group was also kept without HgCl₂.

In respect of the preparation of the stock and test solutions of mercury the test chemical used for the experiment was anhydrous mercury chloride. Mercuric Chloride powder (HgCl₂ =271.50; minimum assay Hg: 98%). The chloride form of the metal was chosen because of its lower toxicity compared to the other forms of mercury [10,11]. After a range finding test, the concentrations prepared for the experiment were 0 mg/L, 0.3 mg/L, 0.5 mg/L, 0.8 mg/L, 1.5mg/L and 3 mg/L of mercury chloride (HgCl₂) respectively. A stock solution of 100mg/L (0.1g/l) of the mercury was prepared by adding 100 mg of mercury to 1 liter of distilled water. Twenty four hours (24) prior to the exposure to mercury chloride and during the whole period of exposure of 96 hours the feeding of the fish was stopped. A static renewal bioassay technique in which the test media were renewed at the same concentration once every twenty four hours was adopted [12]. The bowls were covered with mosquito mesh-sized nets to prevent fish from jumping out or move from one bowl into another. The set up was monitored hourly to observe changes in fish behavior and remove dead fish. A fish was considered dead when observed to be totally immobile with no opercula movement seen when probed with a glass rod.

With respect to the water used in the experiment, its physico-chemical parameters were measured after exposure of the tap water to air in order to lose chlorine. These parameters included temperature, dissolved oxygen, and the potential of hydrogen (pH). They were measured using Horiba


multi-parameter by submersing the sensor in the water at a depth of 10 cm from the water surface. All parameters displaced on the screen of the apparatus.

RESULTS AND DISCUSSION

Bioassay is a necessity to determine the concentration of a toxicant, which could be allowed in waters without adverse effects on the living organisms[14,15]. The physico-chemical parameters of the water measured for were temperature, dissolved oxygen and pH. .

Exposure concentrations of mercury chloride (mg/L)

The physico-chemical parameters recorded were within the permissive limits fixed by WHO which are 6.5 and 6.5- 8.0 respectively for dissolved oxygen and pH. The results of mortality of *Clarias gariepinus* after 96 hours of exposure to mercury chloride are shown in graph 1. It was observed that the mortality recorded in this investigation increased with the rise in concentration. The first death was noticed thirty minutes after the introduction of toxicant in the bowl with the highest concentration in mercury chloride (3mg/L). [16] reported the first death three hours after introduction of toxicant in the exposure of *Clarias gariepinus* to lethal and sub-lethal concentrations of copper. . [17] recorded the first death 36 hours after the exposure to acute toxicity treatments of *Clarias gariepinus* with Synthetic Pyrethroid Deltamethrin, *Parkia biglobosa* and *Raphia vinifera* extracts and Formalin respectively.[18] remarked the first death after thirty hours while treating *Clarias gariepinus* with 120 mg/L of cadmium sulphate.



The duration of resistance of *Clarias gariepinus* in the present study appeared to be the lowest compared to those in the aforementioned studies. Although *Clarias gariepinus* has proved to be very resistant to various toxicants [6, 13], [16-18], it has shown very little resistance to mercury. In the tanks with 2.5mg/L, 3 mg/L & 3.5mg/l concentrations of mercury all the fish died. However no death was recorded in the control tank devoid of mercury. This observation confirmed that the mortality registered could entirely be attributable to the chronic effects of mercury. The estimation of the lethal concentration values (LC 50) was carried out using the linear regression of Probit program. The result is shown in graph-1.

From the graph 1 the 96-hour LC50 value was determined to be 0.60 mg/L with lower and upper confidence limits of 0.135 mg/L and 3.519 mg/L respectively at 95%. The LC50 found in this investigation is similar to that of [19] in the exposure of *Aequidens portalegrensis* to acute concentrations of mercury chloride. [20] reported LC50 of 0.200 mg/L in the treatment of *Poecilia reticulata* with mercury chloride. [21] found LC50 values of 0.168 mg/L and 0.161 mg/L in the exposure of *Variocorhinus barbatulus* and *Zacco barbata* respectively. [22] Recorded 0.22mg/L in an acute mercury toxicity treatment to *Oreochromis niloticus*. The median lethal concentration in our study was the highest recorded compared to those reported in the aforementioned investigations. The chemical product being the same, the difference in the results could be attributable to the

variety in species used. Here, *Clarias gariepinus* proved to be more resistant to mercury chloride than the various species involved in the studies already mentioned.

However the LC50 found in the present study was by far lower than those reported with *Clarias gariepinus* by [23-25]. The difference might be due not only to the various substances and compounds used in the experiments but also the distinct environmental conditions.

Behavioral changes were observed in the catfish in the poisonous solutions. Those symptoms were hyper-activity and attempts to jump out due to skin irritation, restlessness, respiratory distress, loss of balance, gulping for air due to respiratory rate impairment, darkening of the body, sudden and quick movement, rolling movement, back stroke, excessive accumulation of mucus, all these ending in death. The reaction to the toxicant was more noticeable in the media containing the highest two concentrations of mercury chloride.

These observations accord with those remarked by [18-19][24-27] during acute toxicity tests. The accumulation of mucus on the body could be connected to the intensification of mucus secretion of mucous cells activated by the toxicants [28,29]. The ensuing death might be due to increased heart failure, hypertension, gastric hemorrhage, convulsion, paralysis, heart failure and suffocation [30].

CONCLUSION

The investigation of exposing *Clarias gariepinus* to mercury highlighted the high toxicity of mercury by the mortality recorded in the poisonous fish and hence a particular attention should be given to the use of products containing mercury. This study also demonstrated the necessity to regulate the discharge of mercury in effluents from domestic and industrial sources into aquatic systems. There is a need for further research on this subject so as to

establish standards for tropical fish such as *C.gariepinus* and other fish meant for human consumption in throughout Africa in general and particularly in Benin.

REFERENCES

- [1] Bhattacharya T., Chakraborty S., Fadadu B. and Bhattacharya Priya, Heavy metal concentrations in street and leaf deposited dust in Anand city, India, *Res. J. Chem. Sci.*, Vol. **1(5)**, 61-66, (2011)
- [2] Adamu C.I., Nganje T.N., Ukwang E.E., Ibe K.A. and Neji P., A Study of the Distribution pattern of Heavy metals in surface soils around Arufu Pb-Zn mine, Northeastern Nigeria, Using Factor Analysis, *Res. J. Chem. Sci.*, **1(2)**, 70-80 (2011)
- [3] Babu G. R., Reddy B. M., Ramana N.V. and Sashidhar C., Effect of Heavy metal and Magnesium Sulfate on Properties of Blended Cement Mortar, *Res.J. Chem. Sci.*, **1(7)**, 27-32 (2011)
- [4] Ujowundu C.O., Kalu F.N., Nwaoguikpe R.N., Kalu O.I., Ihejirika C.E., Nwosunjoku E.C., Okechukwu R.I., Biochemical and Physical Characterization of Diesel Petroleum Contaminated Soil in South-eastern Nigeria, *Res.J.Chem.Sci.*, **1(8)**, 57-62, (2011)
- [5] Eruola A.O., Ufoegbune G.C., Eruola A.O. Awomeso J.A. and Abhulimen S.A., Assessment of Cadmium, Lead and Iron in Hand Dug Wells of Ilaro and Aiyetoro, Ogun State, South-Western Nigeria, *Res.J.Chem.Sci*, **1(9)**, 1-5 (2011)
- [6] Olaifa F.E. Olaifa A.K., and Onwude T.E., Lethal and sub-lethal effects of copper to the African catfish (*Clarias Gariepinus*) Juveniles, *African Journal of Biomedical Research*, **7**, 65 -70 (2004)
- [7] Sunderland E.M. and Chmura G.L., An inventory of historical mercury emissions in

Maritime Canada: Implications for present and future contamination, *The Science of the Total Environment*, **256**, 39-57 (2000)

- [8] Angle C.R. and McIntire M.S., Red cell lead, whole blood lead, and red cell enzymes, *Environmental Health Perspectives*, **7**, 133-137 (1974)

- [8] Micryakov V.R. and Lapirova T.B., Influence of salts of some heavy metals on the differential blood count in juvenile *Acipenser baeri*, *J. Ichthyol.*, **37(6)**, 458-462 (1997)

- [9] Odiete W.O., Environmental Physiology of Animals and Pollution. First Edition. Diversified Resources Limited, Lagos (1999)

- [10] UNEP (United Nations Environmental Program), Estimation of the lethal toxicity of pollutants in marine fish and invertebrates, *Reference methods for marine pollution studies*, **43**, 27 (1989)

- [11] ASTM (American Society for Testing of Materials) Method 729-90, Guide for Conducting Acute Toxicity Test with Fishes, Macroinvertebrates and Amphibians, ASTM, Philadelphia, 403-422 (1991)

- [12] Okomoda J., Ayuba V.O. and Omeji S., Hematological Changes of *Clarias gariepinus* (Burchell, 1822) Fingerlings Exposed To Acute Toxicity of Formalin, *PAT* 2010; **6 (1)**, 92-10 (2010)

- [13] Standing Committee of Analysts, Acute Toxicity testing with aquatic organisms. Methods for the Examination of Waters and Associated Materials, Her Majesty's Stationery Office, London. (1981)

- [14] Ward G.S. and Parrish P.R., Manual of methods in aquatic environmental research, Part 6, Toxicity Tests, *FAO Fisheries Technical Paper*, **185** FIRI/T185 (1982)

- [15] Datta M. and Kaviraj A., Acute Toxicity of Synthetic Pyrethroid Deltamethrin to Freshwater

- Catfish *Clarias gariepinus*, *Bull. Environment Toxicol*, **70**, 296-29 (2002)
- [16] Fafioye O.O., Adebisi A.A. and Fagade S.O., Toxicity of *Parkia biglobosa* and *Raphia vinifera* extracts on *Clarias gariepinus* juveniles, *African Journal of Biotechnology*, **3(11)**, 627-630 (2004)
- [17] Guedenon P., Edoth P.A., Hounkpatin A. S.Y., Alimba C. G, Ogunkanmi A., Boko M., *Int. J. Biol. Chem. Sci.* **5(6)**, 2497-2501 (2011)
- [18] Hirt L.M. and Domitrovic, H.A. Toxicidad y respuesta histopatológica en *Aequidens portalegrensis* (Pisces, Cichlidae) expuesto a bicloruro de mercurio en ensayos de toxicidad aguda y subletales. Reunion de comunicaciones scientificas tecnologicas, Universidad Nacional del Nordeste, Corrientes, 1999, *Anais... Tomo IV* (Ciências Veterinárias), 19-22 (1999)
- [19] Slabbert J.L. and Venter E.A., Biological assays for aquatic toxicity testing. *Wat. Sci.Tech.*, **39(10-11)**, 367-373 (1999)
- [20] Shyong W.J. and Chen H.C., Acute toxicity of copper, cadmium, and mercury to the freshwater fish *Varicorhinus barbatulus* and *Zacco barbata*, *Acta Zool, Taiwanica*, **11(1)**, 33-45 (2000)
- [21] Ishikawa N.M., Ranzani-Paiva M.J.T., Lombardi J.V., Acute toxicity of mercury (HgCl₂) to Nile Tilapia, *Oreochromis niloticus*, *B. Inst. Pesca*, São Paulo, **33(1)**, 99 – 104 (2007)
- [22] Ayuba V.O., Ofojekwu P.C., Acute toxicity of the root extract of Jimsons Weed *Datura innoxia* to the African catfish *Clarias gariepinus*, *Journals of Aquatic Sciences*, **17(2)**, 131-133 (2002)
- [23] Ezike C, Ufodike E.B.C., Acute toxicity of petrol to the African catfish *Clarias gariepinus*, *Annals of research in Nigeria*, **6**, 1-4 (2008)
- [24] Lawson E.O., Ndimele P/E., Jimoh A.A., Whenu O.O., Acute Toxicity of Lindane (Gamma Hexachloro Cyclohexane) to African Catfish (*Clarias gariepinus*), *Int. J. Anim. Veter. Adv.*, **3(2)**, 63-68 (2011)
- [25] Oti E.E., Acute Toxicity of cassava mill effluent to the African catfish fingerlings, *Journal of Aquatic Sciences*, **20(2)**, 134-137 (2002)
- [26] Oshode O.A., Bakare A.A., Adeogun A.A., Sowunmi A.A., Ecotoxicological Assessment using *Clarias gariepinus* and Microbial Characterization of Leachate from Municipal Solid Landfill, *International Journal of Environmental Research*, **2(4)**, 391-400 (2008)
- [27] Ofojekwu P.C., Effects of sublethal concentrations of *Datura innoxia* leaf on weight gain in the African catfish, *Clarias gariepinus*, *Journal of Aquatic Sciences*, **20(2)** (2005)
- [28] Omitoyin B.O., Metabolic effects of malachite green on *Clarias gariepinus* juveniles, *Journal of Fisheries and Aquatic Science*, **2**, 422-460 (2007).